

Utilizing Remote Sensing Indices to Evaluate Hydrologic Recovery in the Arroyo Seco Watershed

Alicia M. Kinoshita, Terri S. Hogue, and Jongyoun Kim

University of California, Los Angeles
Civil and Environmental Engineering
Hydrology and Water resources

April 3, 2012

2012 Southwest Wildfire Hydrology and Hazards Workshop
Tucson, Arizona



Fire Studies at UCLA



2003 - Old Fire (San Bernardino Mts.)

2005 - Topanga (Malibu Canyon) and Pines Fire (Arroyo Seco)

2006 - Day Fire (Piru Creek-Pyramid Reservoir)

2009 - Station Fire (Angeles NF, Arroyo Seco)

2010 - Bull and Canyon Fires (Sequoia National Forest)

Water quality (stream, reservoirs, soils): Metals (mercury),
nutrients, sediments

Vegetation recovery and coupled hydrologic response, water yield


Modeling post-fire processes - Flooding, water quality, sediment transport



Motivation

Issues

- Increasing wildfires at wildland-urban interface (WUI)
- Significant post-fire consequences
- Uncertainty in long-term hydrologic response



Cal-Adapt, 2011

Methods


Apply remote sensing products to burned watersheds

- I. Previous post-fire recovery work
- II. Water and energy balance variables
 - Evapotranspiration



Goals

- Improve hydrologic estimations and recovery prediction, especially in ungauged/inaccessible areas
- Model development for post-fire prediction






- Moderate Resolution Imaging Spectroradiometer (MODIS)
- Terra and Aqua Satellites
- Advantages: multiple spectral bands, daily overpass
- Disadvantage: Lower spatial resolution (250m)





MODIS Terra Enhanced Vegetation Indices (EVI)




$$EVI = 2.5 \left[\frac{\rho_{NIR}^* - \rho_{RED}^*}{\rho_{NIR}^* + C_1 \rho_{NIR}^* - C_2 \rho_{BLUE}^* + L} \right]$$

- Temp: 16 days
- Spatial: 250 m
- Spatial EVI patterns
- EVI statistical analysis
- Long-term time series




I. Previous study of hydrology and vegetation recovery

Kinoshita and Hogue, 2011




Discharge is not similar to pre-fire conditions...

Kinoshita and Hogue, 2011



Annual basin average EVI
“recovers” by 2010






Annual discharge (cms)



- Pre-fire outliers – El Nino years
- Discharge not similar by 2010

Burn severity, aspect, and post-fire climate affect recovery

Kinoshita and Hogue, 2011




Current Work: Arroyo Seco (41 km²), Angeles National Forest, California



Glendale CIMIS
weather station



Arroyo Seco Hydrology




Arroyo Seco is 100% burned (~moderate severity) by the Station Fire (2009)



Arroyo Seco Hydrology and Vegetation Response (EVI)



January 17, 2009





January 17, 2010

II. Satellite-based Products for Post-fire Water Budgets

(based on work by Kim and Hogue, 2008; 2012)

Approach: Build upon existing theory and develop remotely-sensed hydrologic and energy balance products for burned (ungauged) application


- Evapotranspiration (AET)
- Soil Moisture



Application: Land cover change assessment (fires, urbanization), model development, forcing, validation, regional water budgets, recovery, etc.



Key Alteration - Evaporation and Transpiration (ET) Processes



UCLA ET Algorithms
- Modified Triangle -
- Modified SEBAL

(Kim and Hogue, 2008; 2012)

Net Radiation Product $R_n = SW\downarrow - SW\uparrow + LW\downarrow - LW\uparrow$

Kim and Hogue, 2008

Net Radiation (250x250m, Daily)

Shortwave Radiation

- **MOD03** (1x1km)
Solar Zenith Angle, Geolocation (Lat, Lon)
- **MOD04** (10x10km)
Aerosol Optical Depth, Angstrom Exponent
- **MOD05** (1x1km) - Water Vapor
- **MOD43** (1x1km) - Albedo

Longwave Radiation

- **MOD06** (5x5km)
Total Ozone, Air/Dew Point Temp
- **MOD11** (1x1km)
Emmissivity, Surface Temp

Ground Heat Flux


- **MOD13Q1 & MYD13Q1** (250x250km) EVI

$$G = f(R_n, NDI)$$

Cloudy Condition

- **MOD06** (1x1km)
Cloud Fraction, Cloud Optical Thickness, Surface Temp

Bisht and Bras, 2010; Kim and Hogue, 2011



CIMIS: 388 m

Range of Arroyo Seco: 431-1877 m

June Gloom

Station Fire

Actual Evapotranspiration (UCLA-Triangle Method)

Net Radiation (250x250m, Daily)

Shortwave Radiation

- **MOD03** (1x1km)
Solar Zenith Angle, Geolocation (Lat, Lon)
- **MOD04** (10x10km)
Aerosol Optical Depth, Angstrom Exponent
- **MOD05** (1x1km) - Water Vapor
- **MOD43** (1x1km) - Albedo

Longwave Radiation

- **MOD06** (5x5km)
Total Ozone, Air/Dew Point Temp
- **MOD11** (1x1km)
Emmissivity, Surface Temp

Ground Heat Flux

- **MOD13Q1 & MYD13Q1**
(250x250km) EVI

$$G = f(R_n, NDI)$$

Cloudy Condition


- **MOD06** (1x1km)
Cloud Fraction, Cloud Optical Thickness, Surface Temp

Bisht and Bras, 2010; Kim and Hogue, 2010

+

Evaporative Fraction

- **MOD13Q1** (EVI)
- **MOD11** (LST)




ET
(250x250m, Daily)

$$ET = EF \times (R_n - G)$$

Kim and Hogue, 2012; Jiang and Islam, 2003

Evapotranspiration (UCLA-Triangle Model)

- MODIS ET estimation using VI-LST Triangle Model (Gillies and Carlson, 1995)




$$EF = \frac{ET}{Rn - G} = 1 - \frac{\rho C_p}{Rn - G} \frac{1}{r_a} \Delta T$$


$$EF = 1 - \frac{(1 - EVI^*) \Delta T}{(1 - EVI^*) \Delta T_i^{max} + EVI^* \Delta T_e}$$

(Jiang and Islam, 2003)


MODIS (MOD11) LST -



MODIS (MOD13) VI (NDVI or EVI) -




Spatial ET
Relative to
pre-fire
[mm]



~ JUL 19

~ NOV 1



2006

Jul: 4.7 mm

Nov: 1.2 mm

2007

Jul: 5.0 mm

Nov: 1.8 mm

2008

Jul: 4.8 mm

Nov: 1.4 mm


2009

Jul: 6.2 mm

Nov: 1.1 mm

MODIS ET Validation – comparison to CIMIS ETo

Weekly Comparison



Annual Totals


Year	CIMIS Eto [mm]	MOD ET [mm]
2006	1095	917
2007	1111	1074
2008	1307	1061
2009	1157	894

- 2009 - Station Fire
- CIMIS ETo has a maximum ~6 mm

2009 MODIS Daily ET vs. Weekly CIMIS ET₀



Glendale CIMIS ET₀ (reference ET) measurements over well-watered, maintained grass (~potential)



Higher resolution ET needed for burn areas

Concluding Remarks

- Initial work of the 2003 Old Fire shows post-fire hydrology is controlled primarily by EVI, slope aspect, burn severity
- Climate has shorter-term impacts
- Remote sensing ET over Arroyo Seco captures seasonal trends
- Developed product provides critical spatial and temporal information

Future work


- Higher spatial resolution (MODIS+Landsat product) integrated (*daily, 30m*)
- Produce soil moisture product for burned systems (AMSR-MODIS; Kim and Hogue, 2012)
- Develop short and long-term water balance for burned areas
- Incorporate key variables controlling recovery in multi-variable model to predict daily discharge in ungauged systems

Questions?

Acknowledgments

UCLA helpers:

Sharon Liu
Karen Chu
Carolyn Chou
Audrey Lee



This work is supported by: USFS grant (#1113810020), NASA Earth System Science (ESS) Fellowship (#NNX07AO53H), and NSF Hydrologic Sciences Program (RAPID EAR #0965236)